

Prediabetes in California: Nearly Half of California Adults on Path to Diabetes

Supplemental Tables

LOS ANGELES SPA

Percent of Adults Estimated to Have Prediabetes by Los Angeles County Service Planning Area and Age, California, 2013-14

Los Angeles County Service Planning Area (SPA)	18-39		40-54		55-69		70+		All	
	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Antelope Valley (SPA 1)	30%	(24.3 - 34.7)	50%	(39.9 - 60.7)	59%	(47.9 - 70.7)	48%	(32.7 - 62.6)	42%	(36.5 - 46.7)
San Fernando (SPA 2)	32%	(29.4 - 34.9)	49%	(44.2 - 53.0)	61%	(57.0 - 65.4)	58%	(51.6 - 65.0)	44%	(42.0 - 46.7)
San Gabriel (SPA 3)	35%	(31.3 - 38.0)	48%	(43.1 - 53.7)	55%	(48.5 - 61.3)	50%	(41.9 - 58.8)	45%	(42.5 - 48.0)
Metro (SPA 4)	36%	(32.3 - 39.1)	50%	(42.9 - 58.0)	50%	(36.4 - 62.9)	60%	(50.6 - 68.5)	43%	(39.8 - 46.5)
West (SPA 5)	29%	(25.1 - 33.5)	50%	(44.6 - 56.3)	60%	(53.7 - 66.3)	55%	(46.8 - 62.3)	46%	(43.0 - 50.0)
South (SPA 6)	36%	(31.6 - 39.8)	49%	(39.4 - 59.1)	60%	(52.1 - 67.3)	53%	(38.1 - 67.5)	45%	(41.3 - 49.4)
East (SPA 7)	35%	(31.0 - 38.5)	50%	(43.6 - 56.4)	57%	(49.6 - 65.4)	54%	(41.2 - 66.4)	44%	(41.0 - 47.6)
South Bay (SPA 8)	31%	(27.2 - 34.2)	44%	(38.0 - 50.1)	55%	(48.8 - 61.7)	62%	(55.3 - 68.0)	44%	(40.9 - 46.9)
Los Angeles County	33%	(31.9 - 34.8)	48%	(45.9 - 50.9)	57%	(54.0 - 59.7)	56%	(52.3 - 59.1)	44%	(43.2 - 45.7)

Source: 2013-14 California Health Interview Survey

Note: Estimates of prediabetes are based on predictive models developed using 2009-2012 NHANES data and applied to CHIS 2013-14 data. Prediabetes estimates include adults with undiagnosed diabetes. Nationally, approximately 3.9 percent of adults have undiagnosed diabetes.

Read the full study:

<http://healthpolicy.ucla.edu/publications/search/pages/detail.aspx?PubID=1472>