

# Health Policy Brief

February 2022

## Gaps in Health Care Access and Health Insurance Among LGBT Populations in California

Susan H. Babey, Joelle Wolstein, Jody L. Herman, and Bianca D.M. Wilson

*“LGBT adults are more likely to experience delays in getting needed health care.”*

**SUMMARY:** This study examined differences in health insurance coverage and health care access by sexual orientation and gender identity among California adults. Based on data from the California Health Interview Survey, the results show that although lesbian, gay, and bisexual women and men had similar or better rates of insurance coverage compared to straight women and men, they were more likely to experience barriers in accessing health care, particularly delays in getting needed health care. In addition, gay men, lesbian women, and bisexual women were more likely than straight men and women to report experiencing

unfair treatment when getting medical care. Transgender adults had higher rates of public insurance coverage than cisgender adults but were not more likely to lack health insurance. However, transgender adults were more likely to experience a number of barriers to care, including being less likely to have preventive care visits, more likely to have difficulty finding primary or specialty care providers, and more likely to experience delays in getting needed health care. These findings highlight the need to identify health care and structural interventions that will improve access to care for sexual and gender minorities.

Lesbian, gay, bisexual, and transgender (LGBT) adults in the United States experience many of the same challenges and barriers to accessing health care as straight and cisgender adults, including lack of insurance and poverty. However, research shows that LGBT populations are more likely to be uninsured, to be living in poverty, and to have disabilities that may impact access to health care.<sup>1</sup> Furthermore, sexual and gender minorities have unique barriers to health care that include experiences of discrimination, lack of competent providers, and barriers to gender-affirming health care.<sup>2</sup>


Nationally, research suggests several differences in health care access by sexual orientation.<sup>3</sup> For example, lesbian women and bisexual men and women are more likely than straight women and men to have unmet

medical needs due to cost in the past year. Other research indicates that transgender adults are more likely than cisgender adults to be uninsured and to experience cost-related barriers to health care.<sup>4</sup> Our previous research in California suggested that lesbian, gay, and bisexual women and men have similar or better rates of insurance coverage compared to straight women and men.<sup>5</sup> Despite this, LGB women and men are more likely to experience delays in getting needed health care. Transgender adults are more likely to experience delays in getting medicine that a doctor has prescribed for them.<sup>6</sup>

This report uses data from the California Health Interview Survey (CHIS) to examine differences in health insurance coverage and health care access by sexual orientation and gender identity. The analyses of sexual

## Exhibit 1

## Current Health Insurance Coverage by Sexual Orientation and Gender, Adults Ages 18 and Older, California, 2017–2020



Source: Combined 2017–2020 California Health Interview Surveys

Note: Analysis does not include adults who did not report male or female as their current gender.

\* Significantly different from “Straight or Heterosexual,” with  $p < 0.05$ .

† Estimate is not statistically reliable.

“Gay men and lesbian women had higher rates of employer-based coverage than straight men or women.”

orientation differences include all gender identities (i.e., both transgender and cisgender adults), and the analyses of gender identity differences include people of all sexual orientations. Findings by sexual orientation combine data from 2017 to 2020. Combining data from these recent years allows for the presentation of findings stratified by gender, which is important because disparities vary across lesbian, gay, and bisexual people who identify as either female or male.<sup>7</sup> Findings by gender identity use data from 2015 to 2020. Combining all the years of available data provides for more reliable estimates for this population. Measures are described in more detail under “Data Source and Methods” at the end of this report.

### Gay Men and Lesbian Women More Likely To Have Employer-Based Coverage; Bisexual Men and Women More Likely To Be Insured With Medi-Cal

In 2019–2020, 3.3% (95% CI=3.1–3.6) of California adults described their sexual orientation as lesbian, gay, or homosexual, and an additional 3.6% (95% CI=3.3–3.9) described themselves as bisexual.

The percentage of adults with no health insurance did not vary significantly by sexual orientation (Exhibit 1). However, having employer-sponsored insurance (ESI) or Medi-Cal did. Gay men (57%) and lesbian women (54%) were more likely to have ESI than heterosexual men (48%) or women (44%).

## Indicators of Access to Health Care by Sexual Orientation and Gender, Adults Ages 18 and Older, California, 2017–2020

## Exhibit 2

| Access Indicator                             | Male |          |                          | Female         |          |                          |
|--|------|----------|--------------------------|----------------|----------|--------------------------|
|  | Gay  | Bisexual | Straight or Heterosexual | Gay or Lesbian | Bisexual | Straight or Heterosexual |
| No Usual Source of Care                      | 13%* | 27%*     | 18%                      | 18%            | 24%*     | 12%                      |
| No Doctor Visit in Past Year                 | 13%* | 25%      | 22%                      | 12%†           | 18%*     | 14%                      |
| No Preventive Care Visit in Past Year        | 25%* | 40%      | 32%                      | 30%            | 34%*     | 23%                      |
| Trouble Finding Primary Care Doctor          | 4%   | 5%†      | 4%                       | 5%             | 9%*      | 5%                       |
| Trouble Finding Specialist                   | 13%  | 18%†     | 10%                      | 13%            | 20%*     | 11%                      |
| Delayed or Did Not Get Needed Health Care    | 18%* | 22%*     | 12%                      | 23%*           | 33%*     | 16%                      |
| Delayed or Did Not Get Prescribed Medication | 12%* | 16%*     | 8%                       | 14%            | 21%*     | 11%                      |

Source: Combined 2017–2020 California Health Interview Surveys

Note: Analysis does not include adults who did not report male or female as their current gender.

\* Significantly different from “Straight or Heterosexual,” with  $p < 0.05$ .

† Estimate is not statistically reliable.

Bisexual men and women (31%) were more likely to have Medi-Cal coverage than other groups. The higher rates of public health insurance coverage among bisexual adults compared to monosexual adults (i.e., heterosexual, gay, or lesbian) likely reflects differences in economic stability between these subgroups. Bisexual adults in the U.S., particularly women, have among the highest rates of poverty.<sup>1,3</sup>

### Sexual Minorities, Especially Bisexual Women, Experience Barriers in Access to Health Care


Several indicators of access to care varied by sexual orientation and gender (Exhibit 2). Having a usual source of health care is important for receiving appropriate and timely care. Among women, the proportion with no usual source of care was twice as high for bisexual adults as for heterosexual adults (24% vs. 12%). Bisexual women were also more likely than heterosexual women to have had no doctor visit (18% vs. 14%) and no preventive care visit in the past year (34% vs. 23%). Differences between lesbian and straight women were not statistically significant.

Among men, gay men had the lowest proportion with no usual source of care (13%), significantly lower than the proportions among straight men (18%) or bisexual men (27%). The same pattern is seen for having a doctor visit or a preventive care visit in the past year: Gay men were less likely to have had no doctor visit (13%) and no preventive care visit (25%) in the past year than straight men (22% and 32%, respectively) or bisexual men (25% and 40%, respectively). Differences between straight men and bisexual men were not statistically significant with regard to doctor visits or preventive care visits. Previous research in California indicated that, among men, there were no differences in having a usual source of health care across sexual orientations.<sup>8</sup> However, unlike this current study, the earlier work did not look at differences between gay and bisexual men. This highlights the need to explore within LGB differences when studying sexual minority disparities in health care access.

“Among women, the proportion with no usual source of care was twice as high for bisexual adults as for heterosexual adults.”

## Exhibit 3

## Ever Experienced Unfair Treatment When Getting Medical Care by Sexual Orientation and Gender, Adults Ages 18 and Older, California, 2015–2017



Source: Combined 2015–2017 California Health Interview Surveys  
 Note: Analysis does not include adults who did not report male or female as their current gender. The question about unfair treatment in medical care was included in CHIS 2015–2017 only.

\* Significantly different from “Straight or Heterosexual,” with  $p < 0.05$ .

“Gay men, lesbian women, and bisexual women were more likely than straight men and women to report experiencing unfair treatment when getting medical care.”

Bisexual women were more likely than straight women to experience difficulty finding a primary care provider (9% vs. 5%) and finding a specialist (20% vs. 11%). Among men, there were no statistically significant differences by sexual orientation in those who experienced difficulty finding a primary care provider or finding a specialist.


Sexual minorities were more likely to experience delays in receiving needed medical care or getting prescription medications (Exhibit 2). Among women, the proportion who experienced delays in needed medical care was higher among lesbians (23%) and bisexuals (33%) than among straight women (16%). Bisexual women (21%) were more likely than lesbians (14%) or straight women (11%) to delay getting a prescription medication. Among men, the proportions who experienced delays in getting needed

medical care were higher among gay men (18%) and bisexuals (22%) than among straight men (12%). Bisexual men (16%) and gay men (12%) were more likely than straight men (8%) to delay getting a prescription medication.

There were no statistically significant differences by sexual orientation in the percentage of people who cited the reasons for delaying care provided by the survey. However, in response to a question about whether they had been “treated unfairly” when getting medical care, sexual minorities were more likely to say that they had been (Exhibit 3). More than 40% of lesbian (44%) and bisexual (45%) women reported being treated unfairly, compared to 32% of straight women. Nearly one-third of gay men (32%) reported being treated unfairly, compared to less than one-quarter (23%) of straight men.

**Current Health Insurance Coverage by Gender Identity, Adults Ages 18 and Older, California, 2015–2020**

**Exhibit 4**



Source: Combined 2015–2020 California Health Interview Surveys

\* Significantly different from “Cisgender,” with  $p < 0.05$ .  
 † Estimate is not statistically reliable.

**Transgender Adults More Likely To Have Medi-Cal Coverage, Less Likely To Have Medicare**

In 2019–2020, approximately 0.8% (95% CI=0.6–1.0) of California adults were transgender. The percentages of transgender and cisgender adults with no health insurance did not differ (Exhibit 4). Transgender and cisgender adults had similar rates of insurance coverage through privately purchased plans and employer-based plans. However, transgender adults were significantly more likely than cisgender adults to be covered by Medi-Cal or other public health insurance (37% vs. 21%). Similar to bisexual adults, transgender people have higher rates of

poverty.<sup>1,3</sup> This difference in economic stability between cisgender and transgender people likely contributes to differences in rates of coverage by public health insurance. Transgender adults were also less likely to be covered by Medicare (9% vs. 20%). The difference in Medicare coverage could be explained, at least in part, by differences in age between transgender and cisgender adults, because California adults identifying as transgender tend to be younger.<sup>9</sup>

**Transgender Adults Experience Several Barriers in Access to Health Care**

Transgender and cisgender adults were similar in regard to having a usual source of health

*“Transgender adults were significantly more likely than cisgender adults to be covered by Medi-Cal.”*

## Exhibit 5

## Indicators of Access to Health Care by Gender Identity, Adults Ages 18 and Older, California, 2015–2020

| Access Indicator   | Transgender | Cisgender |
|--|-------------|-----------|
| No Usual Source of Care                                    | 20%         | 16%       |
| No Doctor Visit Past Year                                  | 19%         | 18%       |
| No Preventive Care Visit Past Year                         | 39%         | 28%*      |
| Trouble Finding Primary Care Doctor                        | 8%          | 4%*       |
| Trouble Finding Specialist                                 | 29%         | 11%*      |
| Delayed or Did Not Get Needed Health Care                  | 33%         | 14%*      |
| Delayed or Did Not Get Prescribed Medication               | 23%         | 10%*      |
| Main Reason for Delaying or Not Getting Needed Health Care |             |           |
| Cost or Lack of Insurance                                  | 36%         | 42%       |
| Insurance Not Accepted or Did Not Cover the Care           | 9%          | 3%*       |
| Transportation Problems                                    | 17%         | 2%*       |
| Could Not Get Appointment                                  | 4%†         | 11%*      |
| Did Not Have Time  | 10%         | 19%*      |
| Forgot or Procrastinated                                   | 5%†         | 4%        |
| Anxiety, Fear, Avoid Medical Care                          | 3%†         | 4%        |
| Other  | 15%         | 15%       |

Source: Combined 2015–2020 California Health Interview Surveys

Note: The “Other” category includes a number of different reasons that were provided by smaller proportions of respondents. The most common “other” responses provided were “did not think serious enough” and “not satisfied with care received.”

\* Statistically significant difference between transgender and cisgender, with  $p < 0.05$ .

† Estimate is not statistically reliable.

“Transgender adults were more likely to experience delays in care.”

care and having no doctor visits in the past year (Exhibit 5). However, transgender adults were significantly more likely to have had no preventive care visit in the past year (39% vs. 28%). Transgender adults were also more likely to report having trouble finding a primary care doctor (8% vs. 4%) and finding a health care specialist (29% vs. 11%). It is possible that the much higher rate of having trouble finding health care specialists could be due, at least in part, to difficulty finding providers offering gender-affirming medical care.<sup>10</sup>

Gender minorities were more likely to experience delays in care (Exhibit 5). Transgender adults were significantly more likely than cisgender adults to delay or not get needed health care (33% vs. 14%) and to delay or not get prescribed medications (23% vs. 10%). When asked about the main reason they had delayed or gone without needed health

care, cisgender adults were more likely to report that they did not have enough time or could not get an appointment. Transgender adults were more likely than cisgender adults to report transportation problems as their main reason for delaying or going without needed care (17% vs. 2%), and they were also more likely to report that their insurance was not accepted or did not cover the care (9% vs. 3%).

When asked if they had ever been treated unfairly when getting medical care, a greater percentage of transgender respondents reported experiencing unfair treatment compared to cisgender respondents (42% vs. 28%). Although this difference is large, it was not a statistically significant difference, possibly due in part to the smaller sample size of transgender adults available in the three cycles of data collection that included this question.



## Conclusions

Sexual and gender minorities in California experience a number of barriers in access to health care, despite having similar or better rates of insurance coverage. Gay men were more likely than straight men to have experienced delays in getting needed health care and prescribed medication, even though they were less likely than straight men to have no usual source of care and to have had no preventive care visit in the past year. Bisexual men were more likely than straight men to have no usual source of care and to have experienced delays in getting needed health care and prescribed medication. Bisexual women, in particular, experienced significant barriers to accessing care relative to straight women, including being more likely to have no usual source of care, to have had no doctor visit or preventive care visit in the past year, to have had trouble finding primary care and specialty care providers, and to have delayed or not received needed health care or prescribed medication.

Transgender adults had higher rates of Medi-Cal coverage than cisgender adults, but they were not more likely to be uninsured. Despite this, transgender adults were more likely to have experienced a number of barriers to care, including being less likely to have had preventive care visits, more likely to have had difficulty finding primary or specialty care providers, and more likely to have experienced delays in getting needed health care.

Gay men, lesbian women, and bisexual women were more likely than straight men and women to report experiencing unfair treatment when getting medical care. A higher percentage of transgender adults than cisgender adults also reported experiencing unfair treatment when getting medical care, although this difference was not statistically significant. Other research suggests that sexual and gender minorities experience unique barriers to accessing health care. These include concerns about discrimination

and negative experiences with providers, such as not being believed, being blamed for a health problem, or having their concerns dismissed.<sup>11</sup> Taken together, these findings suggest that previous negative experiences or discrimination could contribute to some of the barriers experienced by sexual and gender minorities, including the higher rates of delayed care.

In our previous examination of differences in access to care by sexual orientation using CHIS data from 2011 to 2014, we found that gay men were less likely than straight men to have no insurance, whereas there was no difference by sexual orientation among women.<sup>5</sup> In addition, employer-based coverage did not vary by sexual orientation, but bisexual men and women were more likely to be covered by Medi-Cal. In contrast to the previous study, the current study found no differences in the percentages of adults without health insurance by sexual orientation among either men or women. This is likely due in part to declines in uninsured rates across all demographic groups following implementation of the Affordable Care Act (ACA) in 2013. In addition, in the current study, gay men and lesbian women were more likely than straight men and women to have ESI. This could be due in part to increases in the proportions of lesbian and gay adults who are married.<sup>12</sup> Similar to the previous study, the current study found that bisexual men and women have higher rates of Medi-Cal coverage than straight and gay or lesbian men and women. These higher rates of Medi-Cal coverage may reflect differences in income and disability rates.<sup>13</sup> Bisexual men and women have higher rates of poverty and disability than straight men and women.

These findings highlight the need for clinical and structural interventions to improve health care access for sexual and gender minorities. The findings reinforce the importance of examining within LGB differences in health care access when studying sexual minority disparities, but they

*“These findings suggest that previous negative experiences or discrimination could contribute to some of the barriers experienced by sexual and gender minorities, including the higher rates of delayed care.”*



The California Health Interview Survey covers a wide array of health-related topics, including health insurance coverage, health status and behaviors, and access to health care. It is based on interviews conducted continuously throughout the year with respondents from more than 20,000 California households. CHIS interviews were offered in English, Spanish, Chinese (both Mandarin and Cantonese), Vietnamese, Korean, and Tagalog. CHIS is designed with complex survey methods requiring analysts to use complex survey weights in order to provide accurate variance estimates and statistical testing. CHIS is a collaboration of the UCLA Center for Health Policy Research, the California Department of Public Health, the California Department of Health Care Services, and the Public Health Institute. For funders and other information on CHIS, visit [chis.ucla.edu](https://chis.ucla.edu).

also suggest the need for additional research to explain existing disparities as well as similarities in access to health care. Future research should also consider within-group differences for transgender men, transgender women, and nonbinary transgender people.

### Data Source and Methods

The findings in this study are based on data from the California Health Interview Survey (CHIS). All analyses presented in this policy brief incorporate survey weights to account for the complex survey design.

Analyses of sexual orientation differences include all gender identities (i.e., they include both transgender and cisgender adults), and the analyses of gender identity differences include people of all sexual orientations. For analyses by sexual orientation, we combined data from 2017 to 2020 to obtain stable estimates and allow for analyses to be stratified by gender (comparing females and males). Adults ages 18 and older were asked to identify their sexual orientation, using the following question: “Do you think of yourself as straight or heterosexual; as gay, lesbian, or homosexual; or as bisexual?” (N=79,965 straight or heterosexual; 2,477 lesbian, gay, or homosexual; and 2,377 bisexual). Responses to this question were used to examine health and access to care by sexual orientation. Analyses stratified by gender are based on self-reported current gender and exclude adults who did not report female or male as their current gender. To examine differences between transgender and cisgender adults, we combined data from all years in which this information was available, 2015 to 2020. To determine whether respondents were transgender or cisgender, adults were asked two questions: “On your original birth certificate, was your sex assigned as male or female?” and “Do you currently describe yourself as male, female, or transgender?” Adults whose assigned sex at birth differs from their current gender identity or who self-report being transgender are transgender (N=451), and those whose sex assigned at birth is the same as their current gender identity are cisgender (N=127,773).

### Author Information

Susan H. Babey, PhD, is a senior research scientist and co-director of the Chronic Disease Research Program at the UCLA Center for Health Policy Research. Joelle Wolstein, MPP, PhD is a research scientist at the UCLA Center for Health Policy Research. Jody L. Herman, PhD, is the Reid Rasmussen Senior Scholar of Public Policy at the Williams Institute at the UCLA School of Law. Bianca D.M. Wilson, PhD, is the Rabbi Zacky Senior Scholar of Public Policy at the Williams Institute at the UCLA School of Law.

### Funder Information

Support for this policy brief was provided by a grant from The California Endowment.

### Acknowledgments

The authors would like to thank the center’s Communications Department for assistance with production and dissemination; Julian Aviles for statistical support; and Andrew Juhnke and Parneet Ghuman for CHIS data access support. The authors are grateful to the following reviewers for their thoughtful and thorough reviews: Krystal Kittle, PhD, postdoctoral research fellow, University of Nevada-Las Vegas; Paul Simon, MD, MPH, chief science officer, Los Angeles County Department of Public Health; and Sean Tan, MPP, senior public administration analyst, UCLA Center for Health Policy Research.

### Suggested Citation

Babey SH, Wolstein J, Herman JL, Wilson BDM. 2022. *Gaps in Health Care Access and Health Insurance Among LGBT Populations in California*. Los Angeles, CA: UCLA Center for Health Policy Research.



10960 Wilshire Blvd., Suite 1550  
Los Angeles, California 90024



The UCLA Center  
for Health Policy Research  
is part of the  
UCLA Fielding School of Public Health.

**UCLA**  
**FIELDING**  
**SCHOOL OF**  
**PUBLIC HEALTH**

**UCLA** School of Law  
**Williams Institute**

The analyses, interpretations, conclusions, and views expressed in this policy brief are those of the authors and do not necessarily represent the UCLA Center for Health Policy Research, the Regents of the University of California, or collaborating organizations or funders.

**PB2022-2**

Copyright © 2022 by the Regents of the University of California. All Rights Reserved.

Editor-in-Chief: Ninez A. Ponce, PhD

Phone: 310-794-0909  
Fax: 310-794-2686  
Email: [chpr@ucla.edu](mailto:chpr@ucla.edu)  
[healthpolicy.ucla.edu](http://healthpolicy.ucla.edu)

## Endnotes

- 1 Badgett MVL, Choi SK, Wilson BDM. 2019. *LGBT Poverty in the United States: A Study of Differences Between Sexual Orientation and Gender Identity Groups*. Los Angeles, CA: The Williams Institute at UCLA School of Law. <https://williamsinstitute.law.ucla.edu/publications/lgbt-poverty-us/>; Herman JL, Wilson BD, Becker T. *Demographic and Health Characteristics of Transgender Adults in California: Findings From the 2015–2016 California Health Interview Survey*. 2017. Los Angeles, CA: UCLA Center for Health Policy Research.
- 2 Kates J, Ranji U, Beamesderfer A, Salganicoff A, Dawson L. 2018. *Health and Access to Care and Coverage for Lesbian, Gay, Bisexual and Transgender (LGBT) Individuals in the U.S.* Menlo Park, CA: Kaiser Family Foundation.
- 3 Wilson BDM, Gordon AR, Mallory C, Choi SK, Badgett MVL, & LBQ Women's Report Team. 2021. *Health and Socioeconomic Well-Being of LBQ Women in the U.S.* Los Angeles, CA: The Williams Institute at UCLA School of Law; Gonzales G, Przedworski J, Henning-Smith C. 2016. Comparison of Health and Health Risk Factors Between Lesbian, Gay, and Bisexual Adults and Heterosexual Adults in the United States: Results from the National Health Interview Survey. *JAMA Internal Medicine* 176(9):1344–51.
- 4 Koma W, Rae M, Ramaswamy A, Neuman T, Kates J, Dawson, L. 2020. *Demographics, Insurance Coverage, and Access to Care Among Transgender Adults*. Menlo Park, CA: Kaiser Family Foundation.
- 5 Wolstein J, Charles SA, Babey SH, Diamant AL. 2018. *Disparities in Health Care Access and Health Among Lesbians, Gay Men, and Bisexuals in California*. Los Angeles, CA: UCLA Center for Health Policy Research.
- 6 Herman JL, Wilson BDM, Becker T. 2017. *Demographic and Health Characteristics of Transgender Adults in California: Findings From the 2015–2016 California Health Interview Survey*. Los Angeles, CA: The Williams Institute at UCLA School of Law, and UCLA Center for Health Policy Research.
- 7 Gonzales G, Henning-Smith C. 2017. Health Disparities by Sexual Orientation: Results and Implications from the Behavioral Risk Factor Surveillance System. *Journal of Community Health* 42(6):1163–1172.
- 8 Wallace SP, Cochran SD, Durazo M, Ford CL. 2011. The Health of Aging Lesbian, Gay, and Bisexual Adults in California. Los Angeles, CA: UCLA Center for Health Policy Research.
- 9 Although mean ages of transgender and cisgender adults were similar in our prior report, that analysis was limited to adults ages 18–70. In our current analysis of California adults ages 18+, the average age of transgender adults was 46, and the average age of cisgender adults was 56.
- 10 James SE, Herman JL, Rankin S, Keisling M, Mottet L, Anafi M. 2016. *The Report of the 2015 U.S. Transgender Survey*. Washington, DC: National Center for Transgender Equality.
- 11 Dawson L, Frederiksen B, Long M, Ranji U, Kates J. 2021. *LGBT+ People's Health and Experiences Accessing Care*. Menlo Park, CA: Kaiser Family Foundation.
- 12 In 2017–20, 26% of homosexual adults ages 18–70 were married, compared to only 13% in 2011–14. Los Angeles, CA: UCLA Center for Health Policy Research. AskCHIS 2011–2014 and 2017–2020. Marital Status by Sexual Orientation, Ages 18–70. Available at <https://ask.chis.ucla.edu/>. Exported on January 14, 2022.
- 13 Fredriksen-Goldsen KI, Kim HJ, Barkan SE. 2012. Disability Among Lesbian, Gay, and Bisexual Adults: Disparities in Prevalence and Risk. *American Journal of Public Health* 102(1):e16–21; LGBT Demographic Data Interactive. January 2019. Los Angeles, CA: The Williams Institute at UCLA School of Law.

